

## NOTE XXXI.

ON THE GENUS CEROBATES SCHH.  
AND DESCRIPTION OF SOME NEW SPECIES

BY

**Dr. ANGELO SENNA,**

Assistant in the R. Museum at Florence.

---

The genus *Cerobates* Schh., widely spread in the oriental, australian and ethiopic regions, is chiefly characterized by having the anterior tibiae notched and strongly toothed on the inner edge, and by having the antennae filiform, not clubshaped; these two characters distinguish it respectively from *Trachelizus* Schh. and *Stereodermus* Lac.

In Lacordaire's classification of the Brenthidæ<sup>1)</sup> *Cerobates* belongs to the group Trachelizides, but in the new arrangement recently proposed by Prof. Sharp in the *Biologia Centrali Americana*<sup>2)</sup>, the Trachelizides are divided into two groups: Stereodermina and Trachelizina. In the former, *Stereodermus*, which has the anterior tibiae more or less notched and the hind coxae more than usually distant from one another, is considered as the typical genus of the group, whilst, in the latter, the typical genus *Trachelizus* has the anterior tibiae unnotched and the hind coxae separated by only the width usual in the Brenthidæ. *Cerobates* should be placed in the Stereodermina.

Lacordaire's group of the Ephemocerides (not »tribe'' as Prof. Sharp writes) is correctly suppressed being a group

---

1) *Genera des Coléoptères*, vol. VII, p. 417, 1866.

2) *Coleoptera*, vol. IV, part 6, p. 7, 1895.

established on two characters — namely the long and slender antennae clothed with delicate hairs, and the large eyes occupying the greater part of the head — remarkably variable in the different genera. *Ephebocerus* Schh. having the anterior tibiae unnotched and being connected with *Trachelizus* by some central-american species is united with the Trachelizina; *Jonthocerus* Lac., which has the inner edge of the anterior tibiae notched and toothed, and which is allied to *Cerobates*, is placed in the Stereodermina. The other genera of the Ephebocerides known at present are: *Hyperephanus* Senna and *Anchisteus* Kolbe: the first has been described by Prof. Sharp under the name of *Homæolizus* and is placed in the Stereodermina, the latter, having the anterior tibiae unnotched, must be united with the Trachelizina. In consequence the Stereodermina are represented by *Stereodermus* Lac., *Stereobates* Sharp, *Hyperephanus* Senna (*Homæolizus* Sharp), *Jonthocerus* Lac. and *Cerobates* Schh. In all these genera the anterior tibiae are constantly more or less notched or toothed on the inner edge, whilst the hind coxae, which are more than usually distant from one another in the typical genus *Stereodermus* Lac., are less widely separated in other genera; on the other hand the distance between these parts in the Trachelizina is also different, so that the only character which satisfactorily distinguishes the two groups is the first mentioned.

The genera of the subfamily Stereoderminae <sup>1)</sup> may be differentiated as follows:

---

1) Prof. Sharp (l. c.) calls this division „group”, not „subfamily” as I do. In his arrangement, the family Brentthidae is divided into two subfamilies: Brentthinae and Ulocerinae, corresponding to Lacordaire’s tribes: Brentthides vrais and Ulocerides. As I retain Ulocerinae does not constitute a subfamily opposite to Brentthinae in Sharp and Lacordaire’s sense, but simply a division of the same value as what is called group by these authors (see my note: Révision des espèces du genre *Ulocerus* Dalm.). I call these groups „subfamilies” as logical subdivisions of the family Brentthidae, and I indicate as groups the successive subdivisions of subfamilies, thence f. i.: family Brentthidae, subfamily Taphroderinae, group Cyphagogi, group Taphroderi, etc.

- I. Antennarum articuli tres apicales  
clavam plus minusve laxam for-  
mantes, semper longiores articulis  
praecedentibus.
- A. Tarsi breves vel modice elon-  
gati, articulo 1<sup>o</sup> ad maximum  
aeque longo duobus sequentibus  
unitis.
- a. Margines sulci prothoracis sim-  
plices vel elevati et rotundati,  
nunquam cariuulati; elytro-  
rum interstitia 2<sup>um</sup> et 3<sup>um</sup> con-  
spicue intus curvata; tibiae  
normales . . . . . *Stereodermus* Lac.
- aa. Margines sulci prothoracis car-  
rinulati; elytrorum interstitia  
2<sup>um</sup> et 3<sup>um</sup> vix intus curvata;  
tibiae lateraliter compressae,  
utrinque carinulatae . . . . . *Stereobates* Sharp.
- AA. Tarsi valde elongati, graciles,  
articulo 1<sup>o</sup> pedum posticorum  
evidenter longiore duobus arti-  
culis sequentibus unitis.
- b. Corpus setosum; antennarum  
clava valde elongata; mar-  
gines sulci prothoracis cari-  
nulati . . . . . *Hyperrephanus* Senna.  
(*Homacolzus* Sharp).
- II. Antennarum articuli tres apicales  
clavam haud formantes: articulo  
apicali solo interdum longiore quam  
praecedente.
- c. Oculi mediocres, antennae  
multo breviores corporis lon-  
gitudine, articulis breviter  
pilosis . . . . . *Cerobates* Schh.
- cc. Oculi in ♂ magni vel maximi,

antennae graciles, totius corporis longitudine parce breviores, articulis elongatis, cylindricis, longe, finissime pilosis . . . . . *Jonthocerus* Lac.

The species of the genus *Cerobates* Schh. are of small size, of graceful form and often closely allied to one another, therefore the determination is not always easy. As I describe here some new species, I believe it useful to give a table for the determination of all the species known at present <sup>1</sup>).

I. Elytrorum angulus apicalis externus rotundatus et marginatus.

A. Prothorax omnino laevis, dorso planatus vel modice convexus.

a. Elytra juxta suturam basi vel usque ad apicem tristriata, versus latera laevius striata aut substriato-punctulata, raro omnino laevia; stria 3<sup>a</sup> dorsali usque ad apicem prolongata, interdum pone medium obsolete, raro oblitterata.

b. Antennarum articulus apicalis conspicue longior articulo praecedente, subcylindricus, apice acuminatus.

c. Metarostrum et mesorostrum sulcata, elytrorum striae impunctatae.

1. Subnitidus; capite, prothorace, elytrorum basi, regione suturali fasciaque transversa postmediana nigris vel ni-

<sup>1</sup>) *Cerobates aciculatus* Walker from Ceylon (Ann. and Magaz. of Nat. Hist., 3rd ser. III, p. 262, 1859) is unknown to me and the description insufficient. I believe this species does not belong to *Cerobates*.

- gro-brunneis, cœterum ferrugineus . . . . . *sulcatus* Boh.
2. Similis et valde affinis sed nitidior, antennarum articulis medianis leviter longioribus. . . . . *sulcirostris* Thoms.
- cc. Metarostrium leviter foveolatum, elytrorum striae punctulatae.
3. Rufo-testaceus, nitidus, elytris dilutioribus, minus nitidis, sutura nigricante . *usambaricus* n.
- bb. Antennarum articulus apicalis vix longior praecedente, ovato-conicus.
- d. Elytra e tertio apicali usque ad apicem regulariter modiceque angustiora.
- e. Prorostrum longitudine subaequale metarostro.
4. Elytra versus latera distincte lineatim punctulata. *complanatus* n.
5. Elytra versus latera plus minusve striata, nunquam omnino laevia; stria 3 dorsali interdum pone medium angustiore, raro oblitterata. *tristriatus* Lund.
- ee. Prorostrum distincte longius metarostro.
6. Elytra versus latera laevia, nitida . . . . . *laevipennis* n.
- dd. Elytra a medio usque ad apicem longe, conspicue angustiora.
7. . . . . *angustipennis* Senna.
- aa. Elytra juxta suturam basi tristriata, lateribus laevissima vel

laevius striata quam dorso; stria 3<sup>a</sup> dorsali basi vel medium versus semper oblitterata.

8. Minor, rostro sat robusto, capite pone oculos rotundato, elytrorum stria 3<sup>a</sup> dorsali medium versus oblitterata, lateribus laevissimis. *sevsulcatus* Motsch.

9. Major, rostro gracili, capite pone oculos sub-angulato, elytrorum stria 3<sup>a</sup> dorsali basin versus oblitterata, lateribus striatis . . . . *adustus* Senna.

AA. Prothorax longitudinaliter sulcatus vel foveolatus, raro basi tenue canaliculatus nunquam omnino laevis.

f. Elytra juxta suturam tristriata, stria 3<sup>a</sup> dorsalis usque ad apicem prolongata.

g. Prothorax tenue interdum obsolete canaliculatus, canaliculo plerumque antice evanescente.

10. . . . . *debilis* Thoms.

gg. Prothorax distincte sulcatus vel foveolatus.

h. Stria 3<sup>a</sup> dorsalis duplo lata praecedentis, interstitia anguste carinata.

11. . . . . *fossulatus* Motsch.

hh. Stria 3<sup>a</sup> dorsalis vix latior praecedente, interstitia minus angusta.

12. Caput pone oculos longe rotundatum, prothoracis impressio antice abbreviata, elytra juxta sutu-

- ram postice sub-quadristriata. . . . . *Grouvellei* Senna.
13. Caput pone oculos breviter rotundatum, prothoracis sulcus ad apicem extensus, elytra antice et postice tristriata . . . *birmanicus* Senna.
- ff.* Elytra juxta suturam dimidio basali tristriata, stria 3<sup>a</sup> dorsali nunquam apicem attingens.
- i.* Elytra versus latera plus minusve distincte striata nunquam omnino laevia.
14. . . . . *canaliculatus* Motsch.
- ii.* Elytra versus latera laevia, nitida.
- j.* Caput pone oculos acute angulatum, prothorax fovea ovata signatus.
15. . . . . *vitiensis* Fairm.
- jj.* Caput pone oculos subrotundatum, prothorax sulcatus usque ad apicem.
16. Metarostrum sublaeve. . . *australasiae* Fairm.
17. Metarostrum distincte sulcatum. . . . . *sumatranus* Senna.
- II. Elytrorum angulus apicalis externus breviter dentatus.
18. Caput pone oculos rotundatum, prothorax sulcatus usque ad apicem, elytra juxta suturam dimidio basali tristriata, stria 3<sup>a</sup> medium versus oblitterata, latera laevia. . . . . *andamanicus* n.

1. *Cerobates sulcatus* Boh.

Schoenherr, Gen. Curcul. V, p. 488, 1840.

Distinctive characters: Teguments hardly shining; head, rostrum and prothorax black or dull brown, elytra with the base, sutural interstice and a transverse band behind the middle of the same colour or slightly paler, the remaining surface ferruginous; apical joint of the antennae distinctly longer than the preceding one, prothorax not furrowed, elytra striate. — Length variable from 5 to 10 mill. — Not common.

*Hab.* Cape of Good Hope and Port Natal.

2. *C. sulcirostris* Thoms.

Thomson, Archives entom. II, p. 119, 1858.

Distinctive characters: Resembles in all respects *C. sulcatus* Boh., but the teguments are more shining, the median antennal joints more elongate. Colour more or less intense according to the size of the specimens which is variable from 5 to 10 mill. — Common.

*Hab.* Gaboon, Congo and Usambara.

3. *C. usambaricus*, n. sp.

*Rufa-testaceus, nitidus, elytris testaceis, minus nitidis, interstitio suturali nigricante; metarostro leviter foveolato, antennarum articulo apicali distincte longiore quam praecedente, prothorace haud sulcato, nitidissimo, elytris dorso et lateribus striatis.* — Long. 7 mill.

*Hab.* Usambara (Plantation Derema).

One specimen collected by Mr. L. Conradt.

Head longer than the breadth at the base, angular behind the eyes, moderately convex above, obsoletely punctured; rostrum one half longer than the head; metarostrium foveolate, prorostrum broader at the tip. Antennae longer than the head and rostrum taken together, with the



median joints elongate, narrower at the base than anteriorly, the 9<sup>th</sup> and 10<sup>th</sup> slightly shorter than the median ones and almost oval; the apical joint cylindrical with the tip acuminate.

Prothorax oblong, not furrowed, obsoletely punctured, very shining. Elytra as long as  $2\frac{1}{2}$  the prothorax, with the base almost truncate, the shoulders rounded, striate above near the sutural interstice and at the sides, obsoletely punctured, the 3<sup>rd</sup> stria is prolonged down to the tip; the dorsal interstices moderately curved, narrower in the middle than in the remaining portion; apex of the elytra margined and rounded at the sides. Legs regular, anterior tibiae moderately toothed on the inner edge.

Body beneath rufous, shining; head and metarostrum at the sides with a line of hair-bearing punctures; prosternum foveolate anteriorly, impressed near the base; metasternum deeply furrowed, abdomen excavated at the base.

#### 4. *C. complanatus*, n. sp.

*C. complanatus* Pow. in litt.

*Depressus, ferrugineo-brunneus, nitidus, capite transverso, pone oculos subrotundato, fronte anguste foveolata; metarostro et mesorostro canaliculatis, antennis gracilibus, articulis medianis subovatis, apicali brevi, ovato-conico; prothorace valde dilatato, depresso, nitidissimo; elytris basi et apici juxta suturam tristriatis, medio bistriatis, stria 3<sup>a</sup> dorsali apicem attingente, versus latera lineatim punctulatis.* — Long.  $5\frac{1}{2}$  mill.

*Hab.* Bonjongo (Cameroon).

One specimen (Civic Museum of Genoa).

Head transverse, emarginate at the base, almost rounded at the sides behind the eyes, hardly convex above, front foveolate.

Metarostrum subconical, canaliculate; mesorostrum almost foveolate, prorostrum a little broader at the tip; antennae slender, short, with the median joints subovate, the apical one also short, ovato-conical.

Prothorax ovate, very broad and depressed, shining, very smooth, the apical margin narrow, truncate, notched at the sides.

Elytra as long as twice the prothorax, broad, emarginate at the base, the shoulders rounded; margined and rounded at the apex, tristriate along the sutural interstice at the base and at the tip, bistriate in the middle, the 3<sup>rd</sup> stria (external) prolonged down to the tip; the sides are punctured.

Legs paler, femora compressed, tibiae very short, the anterior ones deeply notched and toothed on the inner edge.

Body beneath reddish brown, shining, slightly scaled at the sides of the prothorax and in the impressions; head angularly notched at the base, mesorostrum bifoveolate laterally, prosternum depressed, almost marginate on the sides, metasternum and base of the abdomen depressed, the former channeled, apical segment of the latter impressed at the sides and in the middle.

Another specimen from the same locality differs in the following characters: head slightly narrower, more depressed, not foveolate between the eyes; rostrum shorter and broader, more depressed, very obsoletely channeled when viewed sideways; prothorax broader at the apical margin, truncate and less notched at the sides above, elytra shorter and hardly wider; head and metarostrum explanate beneath, the base of the former hairy; mesorostrum impressed, with the margins raised; metasternum with the sides of the median channel oblique inwardly; base of the abdomen broadly excavated, apical segment with the margin of the median foveola raised and hairy. These differences are probably sexual, but I am uncertain whether this specimen is a male or a female.

##### 5. *C. tristriatus* Lund.

*C. Lorquini* Jekel in litt., *Trachelizus pumilus* Dej., *pygmaeus* olim in Dejean's Catalogue.

Lund, Skrivt. af naturhist. selskab. V, 2, p. 66, 1802.

Distinctive characters: Head short, behind the

Notes from the Leyden Museum, Vol. XVII.

eyes almost angulate, never regularly rounded, antennae with the 9<sup>th</sup> and 10<sup>th</sup> joints oval, the apical one short, ovato-conical; prothorax smooth, elytra along the sutural interstice tristriate, the 3<sup>rd</sup> dorsal stria in the typical form prolonged down to the tip; sides of the elytra distinctly striate. Colour ferruginous brown or chestnut, the sutural interstice darker. Sometimes the 3<sup>rd</sup> dorsal stria is narrower or obsolete behind the middle of the elytra. A smaller race (*Trachelizus pumilus* Dej., *pygmaeus olim*) is often pale brick coloured, has the 3<sup>rd</sup> stria interrupted or narrower behind the middle, and the sides of the elytra less evidently striate but never smooth and shining. The passages between the above mentioned forms are almost imperceptible among a great number of specimens and render a satisfactory separation of them very difficult. — Length  $4\frac{1}{2}$ — $8\frac{1}{2}$  mill. — Very common.

*Hab.* From Ceylon to New-Guinea.

6. *C. laevipennis*, n. sp.

*Ferrugineo-brunneus, nitidus, capite brevi, pone oculos subangulato, rostro gracili, longiusculo, parte antica distincte longiore quam postica, hac sublaevi; antennarum articulis 9<sup>o</sup> et 10<sup>o</sup> ovatis, apicali brevi, ovato-conico; prothorace haud sulcato, depresso, nitidissimo; elytris juxta suturam tristriatis, stria externa apicem pertinente, lateribus laevibus, nitidis; apice rotundato et marginato.* — Long. 7 mill.

*Hab.* Sangi Islands.

One specimen in Walter Rothschild's Museum at Tring.

This species is without any doubt the representative of *C. tristriatus* in the Sangi Islands, and differs only from that species by having the rostrum slender, the prorostrum a little longer, the sides of the elytra smooth and shining.

7. *C. angustipennis* Senna.

Notes from the Leyden Museum, XVI, p. 182, 1894.

Distinctive characters: Head short, angulate at

Notes from the Leyden Museum, Vol. XVII.

the sides behind the eyes, metarostrum obsoletely foveolate; antennae with the 9<sup>th</sup> and 10<sup>th</sup> joints almost globuliform, the apical one short; prothorax smooth, depressed; elytra elongate, conspicuously attenuate behind the middle towards the tip, tristriate along the sutural interstice, the 3<sup>rd</sup> dorsal stria not interrupted; the sides are also striate. — Length  $5\frac{1}{2}$ — $7\frac{1}{2}$  mill. — Not common.

*Hab.* Java.

8. *C. sexsulcatus* Motsch.

Motschulsky, Etudes entom. VII, p. 95, 1858.

Distinctive characters: Head regularly rounded behind the eyes: metarostrum almost smooth, apical joint of the antennae short, a little longer than the preceding one, prothorax smooth, shining; elytra along the sutural interstice tristriate in the basal half, the 3<sup>rd</sup> stria (external) interrupted towards the middle of the elytra, never prolonged down to the tip; the sides are smooth, very shining; the apex rounded and margined. — Length  $3\frac{1}{2}$ —6 mill. — Very common.

*Hab.* From Ceylon to New-Guinea.

9. *C. adustus* Senna.

Notes from the Leyden Museum, XVI, p. 184, 1894.

Distinctive characters: Body elongate, colour reddish brown or chestnut, sutural region often darker; head angular at the sides, rostrum slender, antennae with the median joints almost globuliform or slightly oval but shorter than in *C. tristriatus* Lund, apical joint short, ovato-conical; prothorax broad, depressed, smooth. Elytra elongate, tristriate along the sutural interstice in the basal third, bistriate in the remaining portion; the 3<sup>rd</sup> stria (external) interrupted near the base, disc of each elytron smooth, sides striate. — Length 9—10 mill. — Not common.

*Hab.* Java.

Notes from the Leyden Museum, Vol. XVII.

10. *C. debilis* Thoms.

Thomson, Arch. entom. II, p. 118, 1858.

Distinctive characters: Elongate, colour chestnut or reddish brown, the sutural region darker; head short, eyes very approximate at the base, metarostrum foveolate, apical joint of the antennae distinctly longer than the preceding one; prothorax depressed, obsoletely channeled in the basal half, sometimes up to the apex, never smooth; elytra tristriate along the sutural interstice, the 3<sup>rd</sup> dorsal stria prolonged down to the apex, sides striate, obsoletely punctured. — Length  $5\frac{1}{2}$ —8 mill. — Common.

*Hab.* Congo, Gaboon and Zanzibar.

11. *C. fossulatus* Motsch.

Motschulsky, Etud. entom. VII, p. 96, 1858.

Distinctive characters: Head sulcate between the eyes, rostrum sulcate, except near the apex, antennae stoutish, prothorax sulcate up to the apex, elytra along the sutural interstice tristriate, the 3<sup>rd</sup> stria prolonged down to the tip, twice as broad as the internal one, interstices raised, keeled. — Length 5—6 mill. — Rare.

*Hab.* Burma.

12. *C. Grouvellei* Senna.

Bull. Soc. Entom. Ital. XXV, III, p. 307, t. II, f. 6, 1893.

Distinctive characters: Head rounded behind the eyes, furrowed on the front; metarostrum and mesorostrum furrowed; apical joint of the antennae moderately elongate; prothorax foveolate or sulcate, the furrow not prolonged up to the apical margin; elytra along the sutural interstice tristriate in the basal half, quadristriate behind the middle or near the apical declivity, sides very obsoletely striate-punctate. — Length 4— $5\frac{1}{2}$  mill. — Not common.

*Hab.* Sumatra and Brisbane.

13. *C. birmanicus* Senna.

Ann. Soc. Entom. Belgique, XXXVIII, p. 364, 1894.

Distinctive characters: Head shorter than in *C. Grouvellei* Senna, rounded behind the eyes which are more approximate at the base, channeled above between the eyes; metarostrum and mesorostrum furrowed; apical joint of the antennae moderately longer than the preceding one; prothorax furrowed up to the apical margin; elytra along the sutural interstice tristriate, the 3<sup>rd</sup> stria prolonged down to the apex, sides of the elytra obsoletely punctured, not shining. — Length 5—6½ mill. — Not rare.

*Hab.* Burma.

14. *C. canaliculatus* Motsch.

Motschulsky, Etud. entom. VII, p. 96, 1858.

Distinctive characters: Head almost angulate at the sides, front and metarostrum more or less distinctly channeled, apical joint of the antennae moderately longer than the preceding one, prothorax furrowed up to the apical margin, elytra along the sutural interstice in the basal half tristriate, the 3<sup>rd</sup> stria (external) interrupted near the middle of the elytra, sides distinctly striate. — Length 4—5 mill. — Rare.

*Hab.* India and Perak.

15. *C. vitiensis* Fairm.

Le Naturaliste, III, p. 422, 1881; Ann. Soc. Entom. France, 6e sér. I, p. 463, 1881.

Distinctive characters: Head behind the eyes angulate, metarostrum and mesorostrum striolate, apical joint of the antennae moderately longer than the preceding one, prothorax foveolate in the basal half, elytra along the sutural interstice tristriate in the basal half, the 3<sup>rd</sup> stria interrupted near the middle of the elytra, sides smooth, shining. — Length 7½ mill. — Not common.

*Hab.* Fiji Islands.

16. *C. australasiae* Fairm.

Ann. Soc. Entom. France, 6e sér. I, p. 463, 1881.

Distinctive characters: Head behind the eyes almost rounded, front obsoletely foveolate; metarostrum almost smooth; apical joint of the antennae moderately longer than the preceding one, prothorax furrowed up to the apical margin, elytra as in the preceding species but the striae less curved at the base. — Length  $4\frac{1}{2}$ —7 mill. — Not common.

*Hab.* Australia.

17. *C. sumatranus* Senna.

Bull. Soc. Entom. Ital. XXV, III, p. 306, t. III, f. 1, 1893.

Distinctive characters: Closely allied to *C. australasiae* Fairm., but the head slightly longer, the eyes less prominent, the front and metarostrum distinctly furrowed. — Length  $4\frac{1}{3}$ — $5\frac{3}{4}$  mill. — Not common.

*Hab.* Sumatra and Timor.

18. *C. andamanicus*, n. sp.

*Elongatus, angustus, depressus, brunneo-rufus, nitidus, pedibus ferrugineo-brunneis capite pone oculos rotundato, metarostro tenue sulcato, prothorace sulcato usque ad apicem, elytris lateribus sub-parallelis, apice singulatim sub-rotundatis, angulo externo breviter retrorsum dentato, dorso juxta suturam dimidio basali tristriatis, stria 3<sup>a</sup> medium versus oblitterata; lateribus laevibus, nitidis.* — Long.  $6\frac{1}{3}$  mill.

*Hab.* Andaman Islands.

A single specimen in my collection.

Head as long as broad, rounded behind the eyes which are rather large, base emarginate in the middle, slightly convex above; metarostrum slightly furrowed, prorostrum not longer, smooth. Antennae stoutish, with the joints ovate elongate, the apical joint moderately longer than

the preceding one, almost cylindrical, pointed at the tip.

Prothorax oblong-ovate, narrower anteriorly than at the base, depressed above, furrowed up to the apical margin.

Elytra elongate, the sides almost parallel, the apex rounded with the suture emarginate and the external angle shortly dentate; depressed above, tristriate near the sutural interstice in the basal half, the 3<sup>rd</sup> stria is interrupted near the middle of the elytra, sideways they are smooth, shining.

Legs regular. Body beneath more reddish, shining; metasternum and base of abdomen furrowed in the middle.

This species is easily distinguishable from all other known at present, by the peculiar character of the elytra, these having the external apical angles toothed; on the other hand it is closely allied to *C. australasiae* Fairm. and *C. sumatranus* Senna.

Florence, January 25<sup>th</sup> 1896.